

NEUROLOGY

Response of cluster headache to psilocybin and LSD
R. Andrew Sewell, John H. Halpern and Harrison G. Pope, Jr
Neurology 2006;66;1920-1922
DOI: 10.1212/01.wnl.0000219761.05466.43

This information is current as of June 26, 2006

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.neurology.org/cgi/content/full/66/12/1920>

Neurology is the official journal of AAN Enterprises, Inc. A bi-monthly publication, it has been published continuously since 1951. Copyright © 2006 by AAN Enterprises, Inc. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.



Response of cluster headache to psilocybin and LSD

Abstract—The authors interviewed 53 cluster headache patients who had used psilocybin or lysergic acid diethylamide (LSD) to treat their condition. Twenty-two of 26 psilocybin users reported that psilocybin aborted attacks; 25 of 48 psilocybin users and 7 of 8 LSD users reported cluster period termination; 18 of 19 psilocybin users and 4 of 5 LSD users reported remission period extension. Research on the effects of psilocybin and LSD on cluster headache may be warranted.

NEUROLOGY 2006;66:1920–1922

R. Andrew Sewell, MD; John H. Halpern, MD; and Harrison G. Pope, Jr., MD

Cluster headache, often considered the most painful of all types of headache,¹ affects predominantly men (0.4% vs 0.08% of women) and typically begins after age 20 years. The disorder is categorized as either *episodic*, occurring for 1-week to 1-year periods, interspersed with pain-free remission periods, or *chronic*, in which the headaches occur constantly for more than a year with no remission longer than 1 month.² Ten percent of episodic cluster headaches ultimately evolve into the chronic form, and these are termed *secondary chronic*. In standard descriptions of cluster headache, an *attack* refers to the actual paroxysm of pain, a *cluster period* refers to a period of time when attacks occur regularly, and a *remission period* refers to a prolonged attack-free interval.³ Oxygen and sumatriptan are the mainstays of acute abortive treatment, whereas verapamil, lithium, corticosteroids, and other neuromodulators can suppress attacks during cluster periods. No medications are known to terminate cluster periods or extend remission periods. The effects of the ergot alkaloid derivative lysergic acid diethylamide (LSD) and the related indolalkylamine psilocybin on cluster headache have not previously been described and may include such properties.

Case series. We were contacted by a 34-year-old man, diagnosed with episodic cluster headache at age 16 years, who reported a complete remission of his cluster periods when he repeatedly used LSD on a recreational basis between ages 22 and

24 years. Cluster periods resumed once he stopped. Based on this experience, he attempted to treat his cluster headache by ingesting psilocybin-containing mushrooms every 3 months and again achieved lasting remission. On three occasions when he missed his scheduled dose, a cluster period reoccurred.

Intrigued by this history, we located—through cluster headache support groups and an Internet-based survey—several hundred people with cluster headache who reported use of psilocybin-containing mushrooms or LSD specifically to treat their disorder, and we administered a standardized questionnaire (available from the authors). The consent form and study were approved by the McLean Hospital institutional review board. We restricted our analysis to the 53 individuals who 1) agreed to be contacted for evaluation by telephone or e-mail and 2) met International Classification of Headache Disorders-2 criteria for cluster headache and allowed us to obtain copies of medical records documenting a diagnosis of cluster headache by an MD or DO. If the medical records did not support the diagnosis, the subject was excluded from further analysis. The final sample included subjects from across the United States as well as Great Britain, The Netherlands, and South Africa. We found no significant differences between men and women on demographic indices or headache features (table 1). Notably, 31 (58%) of the 53 individuals reported that they had never used psilocybin or LSD except to treat their cluster headache, and a further 13 (25%) had used these drugs for recreational purposes only in the remote past during adolescence.

Results are summarized in table 2 and listed in complete form in table E-1 (on the *Neurology* Web site at www.neurology.org). Of the 32 subjects with episodic cluster headache, 19 had used sublingual psilocybin during cluster attacks; 17 found psilocybin to be effective in aborting attacks (defined as ending the attack within 20 minutes). Only one subject had used sublingual LSD for an acute attack, reporting it to be effective. Twenty-nine subjects had used psilocybin prophylactically during a cluster period; 15 (52%) reported that it was effective (defined as causing total cessation of attacks), and a further 12 (41%) reported partial efficacy (defined as attacks decreasing in intensity or frequency but not ceasing). Five of six LSD users reported cluster period termination. Twenty subjects ingested psilocybin during a remission period; 19 reported an extension of their remission period, in that their next expected cluster period was delayed or prevented entirely. Four of five subjects reported similar remission extension with LSD.

Of the 21 subjects with chronic cluster headache, 5 of 7 reported that psilocybin aborted a cluster attack; 10 of 20 reported that psilocybin induced a complete termination of cluster attacks; and a further 8 reported partial efficacy. Of two chronic cluster headache patients who ingested LSD, both at subhallucinogenic doses, one reported no attacks for 10 days, and the other reported none for 2 months. Interestingly, 22 (42%) of the 53 subjects reported partial or complete efficacy (as defined above) from subhallucinogenic doses of psilocybin or LSD.

Discussion. Our results are interesting for three reasons. First, no other medication, to our knowledge, has been reported to terminate a cluster period. Second, unlike other ergot-based medications, which must be taken daily, a single dose of LSD was described as sufficient to induce remission of a clus-

Additional material related to this article can be found on the *Neurology* Web site. Go to www.neurology.org and scroll down the Table of Contents for the June 27 issue to find the title link for this article.

From the Biological Psychiatry Laboratory (J.H.H., H.G.P.) and Clinical Research Laboratory (R.A.S.), Alcohol and Drug Abuse Research Center, McLean Hospital/Harvard Medical School, Belmont, MA.

Funding sources include MAPS of Sarasota, FL (J.H.H., H.G.P.), and NIDA, NIH T32-DA07252 (R.A.S.). No funding source had any role in study design; collection, analysis, or interpretation of data; writing the report; or submission of the manuscript.

Disclosure: The authors report no conflicts of interest.

Received December 20, 2005. Accepted in final form March 10, 2006.

Address correspondence and reprint requests to Dr. R. Andrew Sewell, Oaks Building, ADARC, McLean Hospital, 115 Mill St., Belmont, MA 02478; e-mail: asewell@mclean.harvard.edu

Table 1 Cluster headache characteristics by sex and subtype

Headache type	n	Age, y	Headache features			
			Attack duration, min	Attacks/day at peak	Cluster period duration, wk	Remission period duration, wk
Episodic						
Men	26	45 (8)	97 (66)	5.5 (3.7)	13 (10)	11 (10)
Women	6	45 (11)	66 (34)	6.2 (3.0)	15 (10)	9 (5)
Total	32	45 (8)	91 (60)	5.6 (3.5)	13 (10)	11 (9)
1° Chronic						
Men	6	48 (8)	79 (57)	9.8 (7.4)	NA	NA
Women	1	38 (NA)	90 (NA)	8.0 (NA)		
Total	7	47 (8)	81 (53)	9.6 (6.8)		
2° Chronic						
Men	10	45 (6)	105 (70)	6.9 (3.0)	NA	NA
Women	4	46 (10)	139 (64)	7.5 (1.0)		
Total	14	45 (7)	115 (68)	7.1 (2.5)		

Data are presented as mean (SD).

1° = primary; 2° = secondary; NA = not applicable.

ter period, and psilocybin rarely required more than three doses. Third, given the apparent efficacy of subhallucinogenic doses, these drugs might benefit cluster headache by a mechanism unrelated to their psychoactive effects.

Table 2 Reported efficacy of principal reported treatments for cluster attacks, cluster periods, and remission extension

Medication	Total, n	Effective, n (%)	Partially effective, n (%)	Ineffective, n (%)
Acute treatment				
Oxygen	47	24 (52)	19 (40)	4 (9)
Triptans	45	33 (73)	8 (18)	4 (9)
Psilocybin	26	22 (85)	0 (0)	4 (15)
LSD	2	1 (50)	0 (0)	1 (50)
Prophylactic				
Propranolol	22	0 (0)	2 (9)	20 (91)
Lithium	20	1 (5)	8 (40)	11 (55)
Amitriptyline	25	0 (0)	4 (16)	21 (84)
Verapamil	38	2 (5)	22 (58)	14 (37)
Prednisone	36	15 (45)	5 (14)	15 (42)
Psilocybin	48	25 (52)	18 (37)	3 (6)
LSD	8	7 (88)	0 (0)	1 (12)
Remission extension				
Psilocybin	22 (31)	20 (91)	NA	2 (9)
LSD	5 (7)	4 (80)	NA	1 (20)

Nine additional individuals had taken psilocybin and two additional had taken lysergic acid diethylamide (LSD) purposefully for remission extension but were not yet due for another cluster period at the time of our evaluation; hence, for them, efficacy could not be scored.

Several limitations of this study should be considered. First, it is subject to recall bias, because it relies primarily on participants' retrospective reports. However, 6 participants (11%) provided detailed headache diaries that corroborated their recall. In addition, 3 (6%) of the 53 participants tried psilocybin for the first time subsequent to consenting to participate in the study but before being questioned; 2 reported complete efficacy and 1 reported partial efficacy—a prospective response rate consistent with our retrospective findings.

A second consideration is the possibility of selection bias, in that individuals with a good outcome may have been more likely to participate. Recruitment over the Internet also selects for younger, more educated, and more motivated subjects,⁴ likely leading to increased reported efficacy.

Third, participants were not blind to their treatment, raising the possibility of a placebo response. However, cluster headache is known to respond poorly to placebo; controlled trials have shown a placebo response of 0% to prophylactic medications such as verapamil,⁵ capsaicin,⁶ and melatonin,⁷ and less than 20% to abortive medications such as sumatriptan.⁸ Therefore, it seems unlikely that we would have found more than 50 cases of apparent response to psilocybin or LSD through placebo effects alone.

Our observations must be regarded as preliminary, in that they are unblinded, uncontrolled, and subject to additional limitations as described above. Therefore, our findings almost certainly overestimate the response of cluster headache to psilocybin and LSD and should not be misconstrued as an endorsement of the use of illegal substances for the self-treatment of cluster headache. However, given the high reported efficacy for this notoriously refrac-

tory condition, it is difficult to dismiss this series of cases as entirely artifactual. Further research is warranted.

Acknowledgment

The authors thank Nancy K. Mello, PhD, and Kimberley Lindsey, PhD, for their comments on an earlier version of this manuscript, and Robert Wold, Earth and Fire Erowid, for assistance with data collection.

References

1. Dodick DW, Rozen TD, Goadsby PJ, Silberstein SD. Cluster headache. *Cephalalgia* 2000;20:787–803.
2. Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders. *Cephalalgia* 2004;24 (suppl 1):44–48.
3. Ekbom K. Some remarks on the terminology of cluster headache. *Cephalalgia* 1988;8:59–60.
4. Etter JF, Perneger TV. A comparison of cigarette smokers recruited through the Internet or by mail. *Int J Epidemiol* 2001;30:521–525.
5. Leone M, D'Amico D, Frediani F, et al. Verapamil in the prophylaxis of episodic cluster headache: a double-blind study versus placebo. *Neurology* 2000;54:1382–1385.
6. Marks DR, Rapoport A, Padla D, et al. A double-blind placebo-controlled trial of intranasal capsaicin for cluster headache. *Cephalalgia* 1993;13:114–116.
7. Leone M, D'Amico D, Moschiano F, Fraschini F, Bussone G. Melatonin versus placebo in the prophylaxis of cluster headache: a double-blind pilot study with parallel groups. *Cephalalgia* 1996;16:494–496.
8. van Vliet JA, Bahra A, Martin V, et al. Intranasal sumatriptan in cluster headache: randomized placebo-controlled double-blind study. *Neurology* 2003;60:630–633.

CME

NEUROLOGY ONLINE CME PROGRAM

Earn CME while reading *Neurology*. This program is available only to online *Neurology* subscribers. Simply read the articles marked CME, go to www.neurology.org, and click on CME Information. This will provide all of the information necessary to get started. The American Academy of Neurology (AAN) is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to sponsor continuing medical education for physicians. *Neurology* is planned and produced in accordance with the ACCME Essentials. For more information, contact AAN Member Services at 800-879-1960.

ONLINE SUPPLEMENTAL DATA

Table (E)T-1. Efficacy of psilocybin and LSD. “Episodic/chronic” = the form of headache; “Psilocybin/LSD treated attack” = psilocybin/LSD taken sublingually aborted an acute attack in 20 minutes or less; “period” = degree of decrease in frequency or intensity of headaches, with “100%” referring to termination of cluster period; “remission” = preventive effect on future cluster periods (episodic) or pain-free time (chronic); “Sub-hallucinogenic?” = dose produces at most a mild subjective sense of altered state of consciousness and brighter colors; “Off meds?” = whether conventional medications are now unnecessary or, if still taken, why; “Can’t obtain” = unable to obtain psilocybin or LSD; “Intensity decreased” = psilocybin or LSD decreased the intensity of but did not eliminate attacks; “1^o” = primary; “2^o” = secondary; “Ø” = not tried.

Case #	Age	Sex	Age of onset	Episodic/Chronic	Psilocybin treated			Sub-hallucinogenic?	LSD treated			Sub-hallucinogenic?	Off meds (or why not)?
					attack	period	remission		attack	period	remission		
1	34	Male	16	Episodic	Yes	100%	Effective	Yes	Ø	100%	Effective	No	Yes
2	43	Male	13	Episodic	No	100%	Effective	Yes	Ø	Ineffective	Ineffective	No	Yes
3	51	Male	25	Episodic	Yes	100%	Effective	No	Yes	100%	Effective	No	Can't obtain
4	36	Male	16	Episodic	Yes	100%	Effective	Yes	Ø	100%	Unclear	No	Yes
5	51	Male	25	Episodic	Yes	>25%	Effective	Yes	Ø	100%	Effective	No	Yes
6	40	Male	19	Episodic	Yes	100%	Effective	Yes					Yes
7	36	Male	30	Episodic	Ø	100%	Effective	Yes					Yes
8	39	Male	20	Episodic	Yes	>75%	Ineffective	Yes					Yes
9	51	Male	48	Episodic	Yes	100%	Not tried	Yes					Yes
10	35	Male	29	Episodic	Yes	100%	Unclear	Yes					Yes
11	40	Male	34	Episodic	Yes	>75%	Not tried	Yes					On abortive only
12	51	Male	46	Episodic	Yes	>75%	Not tried	Yes					On abortive only
13	50	Male	35	Episodic	Ø	>75%	Effective	No	Ø	100%	Unclear	No	Yes
14	45	Male	23	Episodic	Yes	>75%	Effective	No					Yes
15	55	Male	46	Episodic	Ø	100%	Effective	No					Yes
16	34	Male	21	Episodic	Ø	>75%	Effective	No					Yes
17	52	Male	20	Episodic	Ø	>50%	Effective	No					Yes
18	40	Male	16	Episodic	Yes	100%	Unclear	No					Yes
19	50	Male	49	Episodic	Ø	Ø	Unclear	No					Yes
20	54	Male	20	Episodic	Ø	Ineffective	Unclear	No					Yes
21	53	Male	43	Episodic	Ø	100%	Unclear	No					Yes
22	58	Male	35	Episodic	No	>75%	Unclear	No					Yes
23	57	Male	46	Episodic	Yes	100%	Effective	No					Intensity decreased
24	34	Male	18	Episodic	Yes	100%	Unclear	No					Headaches returned
25	36	Male	12	Episodic	Yes	Ø	Effective	No	Ø	Unclear	Effective	No	On abortive only
26	58	Male	38	1° chronic	Ø	100%	N/A	Yes					Yes
27	42	Male	34	1° chronic	Ø	100%	N/A	No	Ø	N/A	10 days	Yes	Yes
28	55	Male	50	1° chronic	Yes	>75%	N/A	No					Intensity decreased
29	51	Male	38	1° chronic	Ø	>25%	N/A	No					Intensity decreased
30	42	Male	34	1° chronic	Yes	100%	N/A	No					Headaches returned
31	40	Male	38	1° chronic					Ø	N/A	2 months	Yes	Headaches returned
32	47	Male	42	2° chronic	Yes	100%	N/A	Yes					Yes
33	41	Male	20	2° chronic	Ø	100%	N/A	Yes					Yes
34	44	Male	36	2° chronic	Ø	100%	N/A	Yes					Yes
35	44	Male	34	2° chronic	No	>75%	N/A	Yes					Verapamil only
36	56	Male	46	2° chronic	Ø	100%	N/A	Yes					Can't obtain
37	46	Male	8	2° chronic	Ø	>75%	N/A	No					Yes
38	36	Male	25	2° chronic	Ø	100%	N/A	No					Headaches returned
39	36	Male	30	2° chronic	Ø	>75%	N/A	No					Headaches returned
40	48	Male	14	2° chronic	Ø	100%	N/A	No					Can't obtain
41	49	Male	30	2° chronic	Ø	>75%	N/A	No					Can't obtain
42	51	Male	50	Episodic	Ø	100%	Effective	Yes					Yes
43	33	Female	16	Episodic	Ø	>75%	Unclear	Yes					Yes
44	48	Female	16	Episodic	Ø	>50%	Unclear	Yes					Yes
45	61	Female	48	Episodic	Ø	>75%	Unclear	No					Yes (not in period)
46	53	Female	18	Episodic	Yes	Ineffective	Effective	No					Yes
47	38	Female	13	Episodic	Ø	100%	Effective	No					On verapamil only
48	38	Female	32	1° chronic	Yes	>50%	N/A	No					Legality concerns
49	43	Female	21	2° chronic	Yes	>75%	N/A	Yes					Intensity decreased
50	36	Female	28	2° chronic	Ø	100%	N/A	No					Yes
51	46	Female	40	2° chronic	Ø	Ø	N/A	No					Yes
52	59	Female	42	2° chronic	No	Ineffective	Ineffective	No					No
53	37	Female	18	Episodic	Yes	Unclear	Effective	Yes					Yes (not in period)